

GenScript *Taq* DNA Polymerase Manual



Technical Manual No. 0213

Version 20060807

I	Introduction	1
II	Application	1
III	Key Features	1
IV	Ship and Storage	1
V	General PCR Protocol Using <i>Taq</i> DNA Polymerase	1
VI	Order Information	2

I. INTRODUCTION

Taq DNA Polymerase is a thermostable DNA Polymerase isolated from an *E. coli* strain that carries the *Taq* DNA polymerase gene. *Taq* DNA polymerase is the most common polymerase used for PCR* reactions

II. APPLICATIONS

Taq DNA Polymerase can be used in most applications including the following:

- PCR.
- 3' A-tailing of blunt ends.
- Primer extension.
- DNA sequencing.

III. KEY FEATURES

Key features of *Taq* DNA Polymerase:

- Terminal transferase activity. *Taq* DNA Polymerase has terminal transferase activity which results in the addition of a single nucleotide (adenosine) at 3' end of the extension product.
- High-purity. No contamination activity has been detected in standard test reactions.

IV. SHIPPING AND STORAGE

This product is shipped on blue ice. Store the product at -20°C .

V. GENERAL PCR PROTOCOL USING *Taq* DNA POLYMERASE

1. Thaw all the reagents for PCR on ice. **Vortex to mix to remove concentration gradient** and then spin down briefly.
2. Set up 50 μl PCR reaction in a thin-wall PCR tube on ice by the following recipe:
 - 5 μl 10X *Taq* buffer solution containing Mg^{2+} .
 - 1 μl 10 mM dNTP stock
 - 1 μl Forward primer (50 μM)



- 1 μ l Reverse primer (50 μ M)
2 μ l Template (up to 100 ng/ μ l) sterile or filtered H₂O
39.5 μ l sterile or filtered H₂O
0.5 μ l *Taq* polymerase (5 units/ μ l)
3. Program PCR cycler as following and start:
Initial denaturing: 94°C for 3 minutes
- Then 30 cycles of: 94°C for 30 seconds
55°C for 45 seconds
72°C for 60 seconds (about 1 kb/minute)
- Extension: 72°C for 7 minutes
4. When the temperature of PCR cycler reaches 94°C, put PCR reaction tube in and continue the program.
5. Analyze PCR fragments on a agarose or polyacrylamide gel.

Note:

1. This is a basic protocol. One needs to optimize the reagent concentrations, conditions and parameters.
2. This protocol is for PCR cycler with a hot lid. Otherwise, mineral oil needs to be added to prevent evaporation.
3. 5% DMSO, 1M betaine, or both can be included in PCR reaction to improve the results when a GC-rich template is used.

VI. ORDER INFORMATION

Taq DNA Polymerase, Cat. No. E00007

Telephone: 732-885-9188, 732-357-3839
Fax: 732-210-0262, 732-885-5878
E-mail: info@genscript.com

For Research Use Only.

* The PCR process is covered by U. S. Patent numbers 4683195 and 4683202 issued to Cetus and owned by Hoffman-La Roche Inc. GenScript does not encourage or support the unauthorized use of the PCR process. Use of this product is recommended for persons that either have a license to perform PCR or are not required to obtain a license.

GenScript Corporation
120 Centennial Ave., Piscataway, NJ 08854
Tel: 732-885-9188, 732-357-3839
Fax: 732-210-0262, 732-885-5878
E-mail: info@genscript.com

Web: <http://www.genscript.com>