

GAPDH Antibody, mAb, Mouse**Cat. No.:** A01622-40**Size:** 40 µg**Synonyms:** GAPDH antibody, mAb, Mouse;**Description:**

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) is well known as one of the key enzymes involved in glycolysis. It catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The enzyme exists as a tetramer of identical chains. Besides its functioning as a glycolytic enzyme in cytoplasm, recent evidence suggest that mammalian GAPDH is also involved in a great number of intracellular processes such as membrane fusion, microtubule bundling, phosphotransferase activity, nuclear RNA export, DNA replication, and DNA repair. During the last decade a lot of findings appeared concerning the role of GAPDH in different pathologies including prostate cancer progression, programmed neuronal cell death, age-related neuronal diseases, such as Alzheimers and Huntingtons disease.

GenScript **GAPDH Antibody, mAb, Mouse** is produced from the hybridoma resulting from fusion of Sp2/0 myeloma and lymphocytes obtained from mouse immunized with human GAPDH protein from erythrocytes.

Immunogen: Human GAPDH protein from erythrocytes**Host:** Mouse**Conjugation:** Unconjugated**Fusion Partner:**

Spleen cells were fused with SP2/0-Ag14 mouse myeloma cells.

Predicated Band Size:

36 KD

Observed Band Size:**Example**

36 KD

Formulation:

0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide.

Clone: 3B1E9**Ig Subclass:** IgG2a, κ**Specificity:** GenScript **GAPDH Antibody, mAb, Mouse**

detects endogenous levels of human, pig, cow, goat, and fish GAPDH.

Positive Control: Hela, HepG2, COS7, and pig muscle.**Purification:** Protein A affinity column**Applications:**


Working concentrations for specific applications should be determined by the investigator. The appropriate concentrations may be affected by secondary antibody affinity, antigen concentration, the sensitivity of the method of detection, temperature, the length of the incubations, and other factors. The suitability of this antibody for applications other than those listed below has not been determined. The following concentration ranges are recommended starting points for this product.

ELISA: 0.05-0.2 µg/ml**Western blot:** 0.1-1.0 µg/ml**Other Applications:** User optimized**Species Reactivity:** Human, pig, cow, goat and fish. This product has been not tested with other species yet.**Reconstitution:**


Reconstitute the lyophilized powder with deionized water (or equivalent) to an antibody concentration of 0.5 mg/ml.

Storage:

The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or below. Avoid repeated freezing and thawing cycles.



Western blot analysis of cell and tissue lysates using GAPDH Antibody, mAb, Mouse (GenScript, A01622, 1 µg/ml). The signal was developed with IRDye™ 800 Conjugated Goat Anti-Mouse IgG. Predicted Size: 36 KD. Observed Size: 36 KD.



Western blot analysis tissue lysates using GAPDH Antibody, mAb, Mouse (GenScript, A01622, 1 µg/ml). The signal was developed with IRDye™ 800 Conjugated Goat Anti-Mouse IgG. Predicted Size: 36 KD. Observed Size: 36 KD.