

Rev04
 Update: Mar,01,2022

DATASHEET


CTLA-4 Fc Chimera, Human

Cat. No.: Z03373

Product Introduction

Species	Human
Protein Construction	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="background-color: #0056b3; color: white; padding: 5px; text-align: center;"> CTLA-4 (Ala37-Phe162) Accession # P16410 </div> <div style="background-color: #76b82a; color: white; padding: 5px; text-align: center;"> hFc </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px;"> N-term C-term </div>
Purity	> 98% as analyzed by SDS-PAGE
Endotoxin Level	< 0.2 EU/μg of protein by gel clotting method
Biological Activity	<p>Assay #1: Measured by its ability to inhibit IL-2 secretion by co-culturing stimulated Jurkat human acute T cell leukemia cells and CD80 expression CHO stable cell line.</p> <p>Assay #2: Immobilized B7-2(CD86), His, Human (Cat. No.: Z03452) at 2.0 μg/ml (100 μl/well) can bind CTLA-4, hFc, Human.</p>
Expression System	CHO
Apparent Molecular Weight	45~48 kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Examples



GenScript product showed equal activity compared to competitor A when binding with biotin CD80 Fc, Human.

Background

Target Background : Cytotoxic T lymphocyte-associated molecule-4 (CTLA-4) is a cell surface molecule that is closely related to CD28, and a powerful negative regulator of T cell activation. Structurally, CTLA-4 is a member of the Ig superfamily, having a single extracellular V-like domain, homology with CD28; The overall sequence homology between CD28 and CTLA-4 is about 20%, but they share a 27% (murine) to 31% (human) identity at the amino acid level. Inhibitory receptor acting as a major negative regulator of T-cell responses. The affinity of CTLA-4 for its natural B7 family ligands, CD80 and CD86, is considerably stronger than the affinity of their cognate stimulatory coreceptor CD28.

Synonyms : CTLA-4

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.