

Rev02
 Update: Aug,08,2025

DATASHEET

CD200/OX-2, His, Cynomolgus

Cat. No.: Z04109

Product Introduction

Species	Cynomolgus
Protein Construction	<div style="display: flex; align-items: center; justify-content: center;"> <div style="background-color: #0056b3; color: white; padding: 5px; margin-right: 10px;"> CD200/OX-2 (Gln56-Gly257) Accession # A0A2K5TQS2 </div> <div style="background-color: #76b82a; color: white; padding: 5px; margin-left: 10px;">His</div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px;"> N-term C-term </div>
Purity	> 95% as determined by BisTris PAGE > 95% as determined by HPLC
Endotoxin Level	Less than 1EU per µg by the LAL method.
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized CD200/OX-2, His, Cynomolgus at 0.5µg/ml (100µl/well) on the plate can bind Human CD200 R1, hFc Tag. Test result was comparable to standard batch
Expression System	HEK293
Theoretical Molecular Weight	23.66 kDa
Apparent Molecular Weight	Due to glycosylation, the protein migrates to 45-55 kDa based on Bis-Tris PAGE result.
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4).
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Background

Target Background : CD200 and its receptors are highly expressed in the lung, on epithelial cells and leukocytes, and emerging evidence links dysregulation of the CD200 pathway with asthma. Moreover, pharmacological modulation of CD200 receptors was shown to improve clinical and inflammatory outcomes of preclinical asthma models.

Synonyms : CD200; MOX1; MOX2; My033; MRC; OX-2; OX2

For research use only. Not intended for human or animal clinical trials, therapeutic or diagnostic use.

Manufacturer: Nanjing GenScript Biotech Co., Ltd. No. 28Yongxi Road, Jiangning District, Nanjing, Jiangsu, China

GenScript USA, Inc.

860 Centennial Ave. Piscataway, NJ 08854

Tel: 1-732-885-9188

