

Rev02
 Update: Aug,08,2025
DATASHEET

IL-2 R beta/CD122[Biotin], His, Cynomolgus

Cat. No.: Z04372

Product Introduction

Species	Cynomolgus
Protein Construction	<div style="display: flex; align-items: center; justify-content: center;"> <div style="background-color: #0056b3; color: white; padding: 5px; text-align: center;"> IL-2 R beta/CD122 (Ala27-Asp239) Accession # Q38J85 </div> <div style="background-color: #76923c; color: white; padding: 5px; text-align: center; margin-left: 10px;"> His </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px; font-size: small;"> N-term C-term </div>
Conjugate	Biotin
Purity	> 95% as determined by BisTris PAGE > 90% as determined by HPLC
Endotoxin Level	Less than 1EU per µg by the LAL method.
Biological Activity	IL-2 R beta/CD122[Biotin], His, Cynomolgus captured on CM5 Chip via AntiHis Antibody can bind Human IL2, No Tag in SPR assay (Biacore T200). Test result was comparable to standard batch.
Expression System	HEK293
Theoretical Molecular Weight	25.6 kDa
Apparent Molecular Weight	Due to glycosylation, the protein migrates to 40-50 kDa based on Bis-Tris PAGE result.
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4).
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

Background

Target Background : IL-2 R beta is a member of the cytokine receptor superfamily. Human IL-2 R beta cDNA encodes a 551 amino acid residue precursor Type I membrane protein with a 26 residue signal peptide, a 214 residue extracellular region, a 25 residue transmembrane region and a 286 residue cytoplasmic domain. Functional IL-2 receptors can exist in two affinity states on cell surfaces, the high affinity complex consisting of heterotrimers of the alpha, beta, and gamma chains, and the intermediate affinity complex comprising heterodimers of the beta and gamma chains.

Synonyms : CD122; IL-15 R beta; IL-2 R beta; IL-2R subunit beta; IL2RB; P70-75; p75; RP5-1170K4.6; IL-15 R β; IL-2 R β; IL-2R subunit β

For research use only. Not intended for human or animal clinical trials, therapeutic or diagnostic use.

Manufacturer: Nanjing GenScript Biotech Co., Ltd. No. 28Yongxi Road, Jiangning District, Nanjing, Jiangsu, China