


Rev02
 Update: Aug,08,2025
DATASHEET

FGFR3 beta (IIIc), His & Avi, Human

Cat. No.: Z04263

Product Introduction

| | |
|-------------------------------------|--|
| Species | Human |
| Protein Construction |  <small>N-term</small> <small>C-term</small> |
| Purity | > 95% as determined by BisTris PAGE > 95% as determined by HPLC |
| Endotoxin Level | Less than 1EU per µg by the LAL method. |
| Biological Activity | Measured by its binding ability in a functional ELISA. Immobilized FGFR3 beta (IIIc), His & Avi, Human at 0.5µg/ml (100µl/Well) on the plate can bind AntiFGFR3 Antibody, hFc Tag. Test result was comparable to standard batch. |
| Expression System | HEK293 |
| Theoretical Molecular Weight | 30.1 kDa |
| Apparent Molecular Weight | Due to glycosylation, the protein migrates to 50-60 kDa based on Bis-Tris PAGE result. |
| Formulation | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage & Stability | Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles. |

Background

Target Background : Four distinct genes encoding closely related FGF receptors, FGF R1-4, are known. All four genes for FGF Rs encode proteins with an N-terminal signal peptide, three immunoglobulin (Ig)-like domains, an acid-box region containing a run of acidic residues between the Igl and IgII domains, a transmembrane domain and the split tyrosine-kinase domain. FGFR3 is tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of cell proliferation, differentiation and apoptosis. Plays an essential role in the regulation of chondrocyte differentiation, proliferation and apoptosis, and is required for normal skeleton development. Regulates both osteogenesis and postnatal bone mineralization by osteoblasts.

Synonyms : ACH; CD333; CEK; CEK2; EC 2.7.10; FGF R3; FGFR3; HSFGR3EX; JTK4

For research use only. Not intended for human or animal clinical trials, therapeutic or diagnostic use.

Manufacturer: Nanjing GenScript Biotech Co., Ltd. No. 28Yongxi Road, Jiangning District, Nanjing, Jiangsu, China