

Rev02  
 Update: Aug,08,2025
**DATASHEET**

# FGFR3 beta (IIIb)[Biotin], His & Avi, Human

Cat. No.: Z04090

## Product Introduction

<b>Species</b>	Human
<b>Protein Construction</b>	<div style="display: flex; align-items: center; justify-content: center;"> <div style="background-color: #0056b3; color: white; padding: 5px; text-align: center;"> <b>FGFR3 beta (IIIb) (Asp127-Gly377)</b>            Accession # P22607-2         </div> <div style="background-color: #90c080; color: white; padding: 5px; text-align: center; margin: 0 5px;">His</div> <div style="background-color: #008000; color: white; padding: 5px; text-align: center; margin: 0 5px;">Avi</div> </div> <div style="display: flex; justify-content: space-between; width: 100%; margin-top: 5px;"> <span>N-term</span> <span>C-term</span> </div>
<b>Conjugate</b>	Biotin
<b>Purity</b>	> 95% as determined by BisTris PAGE > 95% as determined by HPLC
<b>Endotoxin Level</b>	Less than 1EU per µg by the LAL method.
<b>Biological Activity</b>	Measured by its binding ability in a functional ELISA. Immobilized FGFR3 beta (IIIb) [Biotin], His & Avi, Human at 0.5µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for AntiFGFR3 Antibody, hFc Tag. Test result was comparable to standard batch.
<b>Expression System</b>	HEK293
<b>Theoretical Molecular Weight</b>	30.5 kDa
<b>Apparent Molecular Weight</b>	Due to glycosylation, the protein migrates to 40-55 kDa based on Bis-Tris PAGE result.
<b>Formulation</b>	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4).
<b>Reconstitution</b>	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage &amp; Stability</b>	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

## Background

**Target Background :** Four distinct genes encoding closely related FGF receptors, FGF R1-4, are known. All four genes for FGF Rs encode proteins with an N-terminal signal peptide, three immunoglobulin (Ig)-like domains, an acid-box region containing a run of acidic residues between the IgI and IgII domains, a transmembrane domain and the split tyrosine-kinase domain. FGFR3 is tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of cell proliferation, differentiation and apoptosis.

**Synonyms :** ACH; CD333; CEK; CEK2; EC 2.7.10; FGF R3; FGFR3; HSFGR3EX; JTK4

**For research use only. Not intended for human or animal clinical trials, therapeutic or diagnostic use.**

Manufacturer: Nanjing GenScript Biotech Co., Ltd. No. 28Yongxi Road, Jiangning District, Nanjing, Jiangsu, China