

Rev03
Update: Aug,08,2025**DATASHEET**


CD27 Ligand/CD70, His, Cynomolgus

Cat. No.: Z04112


Product Introduction

Species	Cynomolgus				
Protein Construction	<table border="1"><tr><td>His</td><td>CD27 Ligand/CD70 (Gln39-Pro194) Accession # G7PYU6-1</td></tr><tr><td>N-term</td><td>C-term</td></tr></table>	His	CD27 Ligand/CD70 (Gln39-Pro194) Accession # G7PYU6-1	N-term	C-term
His	CD27 Ligand/CD70 (Gln39-Pro194) Accession # G7PYU6-1				
N-term	C-term				
Purity	> 95% as determined by Bis-Tris PAGE				
Endotoxin Level	Less than 1EU per ug by the LAL method.				
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized CD27 Ligand/CD70, His, Cynomolgus at 2µg/ml (100µl/well) on the plate can bind Cynomolgus/Rhesus macaque CD27, hFc Tag. Test result was comparable to standard batch.				
Expression System	HEK293				
Theoretical Molecular Weight	18.4 kDa				
Apparent Molecular Weight	Due to glycosylation, the protein migrates to 55-70 kDa based on Bis-Tris PAGE result.				
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4).				
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.				
Storage & Stability	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.				

Examples



CD27 Ligand/CD70, His, Cynomolgus on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.



Immobilized CD27 Ligand/CD70, His, Cynomolgus, His Tag at 2 µg/ml (100 µL/well) on the plate. Dose response curve for Cynomolgus / Rhesus macaque CD27, hFc Tag with the EC₅₀ of 100.2 ng/ml determined by ELISA.

Log Cynomolgus / Rhesus macaque CD27, hFc Tag Conc. (µg/ml)

Background

Target Background : CD70, also named CD27 ligand (CD27L), is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and has been designated TNFSF7. CD70 is a cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.

Synonyms : CD70 molecule; CD70; TNFSF7; CD27 Ligand; CD27-L; CD27LG; Ki-24 antigen; TNFSF7G

For research use only. Not intended for human or animal clinical trials, therapeutic or diagnostic use.

Manufacturer: Nanjing GenScript Biotech Co., Ltd. No. 28Yongxi Road, Jiangning District, Nanjing, Jiangsu, China