

Rev05
 Update: Aug,11,2025
DATASHEET

IFN- ω , Human

Cat. No.: Z02909

Product Introduction

Species	Human
Protein Construction	IFN- ω (Cys24-Ser195) Accession # P05000
Purity	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
Endotoxin Level	< 1 EU/ μ g of protein by LAL method
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a chemotaxis bioassay using human TF-1 cells is less than 0.01 ng/ml, corresponding to a specific activity of > 1.0 \times 10 ⁸ IU/mg.
Expression System	E. coli
Theoretical Molecular Weight	20 kDa
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS, pH 7.4.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Background

Target Background : Interferon-Omega (IFN- ω) coded by IFNW1 gene in human, is a member of the type I interferon family, which includes IFN- α , IFN- β , and IFN- ω . The IFNAR-1/IFNAR-2 receptor complex can help with the signal transduction, followed the antiviral or the antiproliferative actions. IFN- ω is derived from IFN- α/β and share 75% sequence with IFN- α . It has two intramolecular disulfide bonds which are crucial for activities. Mire-Sluis et al have described bioassays for IFN- α , IFN- β , and IFN- ω that exploit the ability of these factors to inhibit proliferation of TF-1 cells induced by GM-CSF. The bioassays can be used also with Epo and TF-1 cells, or Epo and Epo-transfected UT-7 cells.

Synonyms : IFN Omega; IFN- ω

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.

GenScript USA, Inc.

860 Centennial Ave. Piscataway, NJ 08854

Tel: 1-732-885-9188

Confidential and Privileged



Manufacturer: Nanjing GenScript Biotech Co., Ltd. No. 28Yongxi Road, Jiangning District, Nanjing, Jiangsu, China

GenScript USA, Inc.

860 Centennial Ave. Piscataway, NJ 08854

Tel: 1-732-885-9188