

Human Recombinant Gα15 Stable Cell Line

Cat. No. M00257

Version 07272020

I. Introduction	1
II. Background	1
III. Thawing and Subculturing.....	2
IV. References.....	3
V. Limited Use License Agreement	4

I. INTRODUCTION

Catalog Number: M00257

Cell Line Name: CHO-K1/Gα15

Gene Synonyms: GNA15, GNA16

Official Full Name: Guanine nucleotide binding protein (G protein), alpha 15 (Gq class)

Expressed Gene: GenBank Accession Number NM_002068; no expressed tags

Host Cell: CHO-K1

Quantity: Two vials of frozen cells (>1×10⁶ per vial)

Application: Functional assay for Gs and Gi/o-coupled GPCR receptors

Freeze Medium: 45% complete growth medium, 45% FBS (Cat. #10099-141, Gibco), 10%

DMSO (Cat. #D2650, Sigma)

Complete Growth Medium: Ham's F-12K (Kaighn's) (Cat. #21127, Life Technologies), 10% FBS

Culture Medium: Ham's F-12K (Kaighn's), 10% FBS, 100 µg/ml Hygromycin B (Cat. #10687010, Invitrogen)

Mycoplasma Status: Negative*

Storage: Liquid nitrogen immediately upon receipt

** The mycoplasma test was performed with MycoAlert™ PLUS Mycoplasma Detection Kit of Lonza.*

II. BACKGROUND

1. CHO-K1/Gα15

CHO-K1/Gα15 is a CHO-K1 cell line stably expressing the Gα15 alpha subunit protein which a Gq protein. It is used as a host cell for transfection expression of Gs and Gi/o -coupled receptors, the constitutively expressed Gα15 protein in the cells allows many transfected receptors which normally stimulate/inhibit the cAMP pathway, to couple to Gq signal transduction and mobilize intracellular calcium. The cell line carries the hygromycin B resistance gene and is resistant to hygromycin B.

2. The sequence of Gα15

ATGGCCCGCTCGCTGACCTGGCGCTGCTGCCCTGGTGCCTGACGGAGGATGAGAAGGC

CGCCGCCCGGGTGGACCAGGAGATCAACAGGATCCTCTTGGAGCAGAAGAAGCAGGACC
GCGGGGAGCTGAAGCTGCTGCTTTTGGGCCAGGCGAGAGCGGGAAGAGCACCTTCATC
AAGCAGATGCGGATCATCCACGGCGCCGGCTACTCGGAGGAGGAGCGCAAGGGCTTCCG
GCCCCTGGTCTACCAGAACATCTTCGTGTCCATGCGGGCCATGATCGAGGCCATGGAGCG
GCTGCAGATTCCATTCAGCAGGCCCGAGAGCAAGCACCACGCTAGCCTGGTCATGAGCCA
GGACCCCTATAAAGTGACCACGTTTGGAGAAGCGCTACGCTGCGGCCATGCAGTGGCTGTG
GAGGGATGCCGGCATCCGGGCCTACTATGAGCGTCGGCGGGAATTCCACCTGCTCGATTC
AGCCGTGTAACCTGTCCCACCTGGAGCGCATCACCGAGGAGGGCTACGTCCCCACAGC
TCAGGACGTGCTCCGCAGCCGCATGCCACCACTGGCATCAACGAGTACTGCTTCTCCGT
GCAGAAAACCAACCTGCGGATCGTGGACGTGCGGGGCCAGAAGTCAGAGCGTAAGAAATG
GATCCATTGTTTCGAGAACGTGATCGCCCTCATCTACCTGGCCTCACTGAGTGAATACGAC
CAGTGCCTGGAGGAGAACAACCAGGAGAACCGCATGAAGGAGAGCCTCGCATTGTTTGGG
ACTATCCTGGAACCTACCCTGGTTCAAAGCACATCCGTCATCCTCTTTCTCAACAAAACCGA
CATCCTGGAGGAGAAAATCCCCACCTCCCACCTGGCTACCTATTTCCCCAGTTTCCAGGGC
CCTAAGCAGGATGCTGAGGCAGCCAAGAGGTTTCATCCTGGACATGTACACGAGGATGTAC
ACCGGGTGCGTGGACGGCCCCGAGGGCAGCAAGAAGGGCGCACGATCCCGACGCCTCTT
CAGCCACTACACATGTGCCACAGACACACAGAACATCCGCAAGGTCTTCAAGGACGTGCG
GACTCGGTGCTCGCCCGCTACCTGGACGAGATCAACCTGCTGTGA

III. THAWING AND SUBCULTURING

Thawing protocol

1. Remove the vial from liquid nitrogen tank and thaw cells quickly in a 37°C water-bath.
2. Just before the cells are completely thawed, decontaminate the outside of the vial with 70% ethanol and transfer the cells to a 15 ml centrifuge tube containing 9 ml of complete growth medium.
3. Pellet cells by centrifugation at 200 x g force for 5 min, and discard the medium.
4. Resuspend the cells in complete growth medium.
5. Add 10 ml of the cell suspension in a 10 cm dish.
6. Add Hygromycin B to a concentration of 100 µg/ml in the following day.

Subculturing protocol

1. Remove and discard culture medium.
2. Wash cells with PBS (pH=7.4) to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 ml of 0.05% (w/v) Trypsin- EDTA (GIBCO, Cat No. 25300) solution to 10 cm dish and observe the cells under an inverted microscope until cell layer is detached (usually within 3 to 5 minutes).

Note: To avoid clumping, do not agitate the cells by hitting or shaking the dish while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate detaching.

4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting, centrifuge the cells 200 x g force for 5min, and discard the medium.
5. Resuspend the cells in the culture medium and add appropriate aliquots of the cell

suspension to new culture vessels.

6. Incubate cultures at 37°C.

Subcultivation Ratio: 1:3 to 1:8

Medium Renewal: Every 2 to 3 days

IV. REFERENCES

1. Elmenhorst, D., Meyer, PT., Winz, OH., *et al.* (2007) Sleep deprivation increases A1 adenosine receptor binding in the human brain: a positron emission tomography study. *J. Neurosci.* 27 (9): 2410–2415
2. Fredholm, BB., IJzerman, AP., Jacobson, KA., *et al.* (2001) International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol. Rev.* 53 (4): 527–552.

GenScript USA Inc,
860 Centennial Ave.
Piscataway, NJ 08854
Toll-Free: 1-877-436-7274
Tel: 1-732-885-9188, Fax: 1-732-210-0262
Email: product@genscript.com
Web: <http://www.genscript.com>
For Research Use Only.

Limited Use License Agreement

This is a legal agreement between you (Licensee) and GenScript USA Inc. governing use of GenScript's stable cell line products and protocols provided to licensee. By purchasing and using the stable cell line, the buyer agrees to comply with the following terms and conditions of this label license and recognizes and agrees to such restrictions:

- 1) The products are not transferable and will be used at the site where they were purchased. Transfer to another site owned by buyer will be permitted only upon written request by buyer followed by subsequent written approval by GenScript.
- 2) The purchaser cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party.
- 3) The products sold by GenScript are for laboratory and animal research purposes only. The products are not to be used on humans, for consumption, or for any unlawful uses.

GenScript USA Inc. will not assert against the buyer a claim of infringement of patents owned or controlled by GenScript USA Inc. and claiming this product based upon the manufacture, use or sale of a clinical diagnostic, therapeutic and vaccine, or prophylactic product developed in research by the buyer in which this product or its components has been employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on the use of this product for other purposes, contact Marketing Department, GenScript USA Inc., 860 Centennial Ave. Piscataway, NJ 08854, U.S.A. Phone: 1-732-885-9188. Fax: 1-732-210-0262. Email: marketing@genscript.com.